



ieMR Advanced

Continuous infusions – begin insulin infusion

Medication Management

Quick reference guide

Begin Insulin Infusion using Interactive View

1. Navigate to the **Interactive View** from the menu.
2. Click on Paediatric Quick View and navigate to the vital signs tab.
3. Enter the Patients BGL and verify by clicking the green tick.
4. Perform PPID and allergy check, mark allergies as reviewed Mark All as Reviewed
5. Navigate to the **MAR** and view the order, hover to see special instructions and prepare the medication.
6. Navigate to the **MAW**, open the administration window by clicking on the results column of the infusion order. The Default action will be 'Begin Bag'.

The **Administration** window will open.

7. Select ****Peripheral Intravenous Cannula** from the ***Site** dropdown menu.

8. To check the **Target BGL range as per prescriber**, right click on the **Order sentence** to view the **Order Info**.

9. Close the **Order Info**.

10. Based on the order info enter the insulin dose required – Units/kg/hr.

- 11.

Note: The ***Rate** (ml/hr) will autopopulate.



Note: The *Insulin dose* is the strength/dose that the patient will be receiving. The Rate is what the infusion is running at, measured in mL/hr.

12. Enter the nurse witness.
13. Click *OK* to save the changes.
14. You will be prompted for the *Nurse Witness's Password*.
15. Click *OK*.
16. Sign by clicking on the *Sign* in the bottom right of the *MAW* pop up box

Note: The *Diabetes – Endocrine iView band* will populate information regarding insulin, BGLs and carbohydrate intake.

