



ieMR Advanced

Continuous infusions: modify


Quick reference guide

To modify any infusion result, follow the steps below:


1. Click **MAR** from the **Menu**.
2. Right click the Administration Information IV event you need to modify.
3. Select **Modify**.




The **Administration** window will open.




4. Make the modifications as required.
5. Click **Comment...** to provide the modification reason and click **OK**.



6. Click **Apply**.



Note: Icon  shows a modified comment.

7. Click the  to sign.
8. **Refresh** the page.



PRN			
oxycodone 5 mg, Tablet, Oral, 4 hourly, start: 08/17/16 4:16:00 AEST, PRN for pain/fever	PRN	5 mg Not given within 7 days.	
oxycodone Pain Response			
Continuous Infusions			
heparin additive 25,000 unit(s) [1,000 u... sodium chloride 0.9% PREFILLED SYRINGING... Rate: 2 ml/hr, Total volume (ml): 50, 50 ml, IV continuous infusion, start: 08/25/16 5:33:00 AEST, Indication: Acute Treatment of Deep Vein Thrombosis or Pulmonary Embolism, Initial max rate 1,500 units/hr		Pending Not given within 7 days.	
Administration Information			
heparin Sodium Chloride 0.9% IV			
Sodium Chloride 0.9% intravenous infusi... Rate: 100 ml/hr, Total volume (ml): 1,000, 1,000 ml, Solution, injectable, IV continuous infusion, start: 10/06/16 7:24:00 AEST, stop: 10/06/16 17:23:00 AEST, order duration: 1 ba...		Pending not bag started: 26-Oct-2016 9:25 AEST	End Bag 1 of 1
Administration Information			
Sodium Chloride 0.9% IV	100 ml, Vivast 100 ml, Avth		* (c) 60 ml, Waste No Site Change, Beon Bag 1,1 * (c) 60 ml, Modified

Note: The *(c) symbol will display on the relevant box to indicate that there has been a change and a comment has been made against it.

