



Ketamine Infusions - Setting up a Dynamic Group

Medication Management

Quick reference guide

This should be completed when commencing the infusion.

- In the MAR, R) click on the Ketamine order and select **Order Information**. Select **Tear Off** from the toolbar to view the order in a second window. Position this so you can read the order.
 
- Open **Interactive View** in the main window.
- Navigate to the **Paediatric Lines and Devices** band and select the **Pain Infusion** subheading.
 
- Select the rubicks cube icon to create a label for the dynamic group for the infusion.

Pain Infusion



- Select **Drug infusing** – Ketamine - then **OK**.

Drug Infusing:

<input type="checkbox"/>	Fentanyl
<input type="checkbox"/>	Morphine
<input type="checkbox"/>	Oxycodone
<input type="checkbox"/>	Hydromorphone
<input type="checkbox"/>	Buprenorphine
<input checked="" type="checkbox"/>	Ketamine
<input type="checkbox"/>	Ropivacaine
<input type="checkbox"/>	Bupivacaine
<input type="checkbox"/>	Levo-Bupivacaine

- Complete the following:
 - Is this a maternity patient** – No.
 - Assessment type** – Routine.
 - Infusion type** - PCA/NCA/MCA/Continuous opioid infusion.

- PCA mode** – Continuous infusion.
- Number of drugs infusing can be left blank as drug name is included in the dynamic group label.
- Continuous infusion rate** – enter rate.
- Continuous infusion unit** – mL/hr.
- Line labelled and checked** – Yes.
- Nurse/Midwife Witness** – Yes.

◀Ketamine▶	
◊ Is this a Maternity patient?	No
◊ Assessment type	Routine
◊ Activity	
◊ Infusion type	PCA/NCA/MCA/Continuous opioid infusion
◊ Number of drugs infusing	
◊ PCA mode	Continuous infusion
◊ Pump settings checked & locked	Yes
◊ Continuous infusion rate	1
◊ Continuous infusion unit	mL/hr
Total cumulative dose	
Total cumulative dose unit	
24 hour cumulative dose	
24 hour cumulative dose unit	
◊ Line labelled and checked	Yes
◊ Patent IV access	
◊ Clinician bolus dose given	
Nurse/Midwife Witness	Yes

Documenting at the Change of Shift

- Navigate to the **Continuous Infusions** section of the **MAR**.
- Review the order.
- Navigate to **iView** and locate the dynamic group for the infusion.
- In assessment type enter **Change of caregiver**
- Complete the following fields:
 - Infusion type** – PCA/NCA/MCA/Continuous opioid infusion.
 - Continuous infusion rate** – enter rate.
 - Continuous infusion unit** – mL/hr.
 - Line labelled and checked** – Yes.
 - Nurse/Midwife Witness** – Yes.



ieMR Advanced

▲ <Ketamine>	
◆ Is this a Maternity patient?	
◆ Assessment type	Change of care giver
◆ Activity	
◆ Infusion type	PCA/NCA/MCA/Continuous opioid infusion
◆ Number of drugs infusing	
◆ PCA mode	Continuous infusion
Pump settings checked & locked	Yes
◆ Continuous infusion rate	1
◆ Continuous infusion unit	mL/hr
Total cumulative dose	
Total cumulative dose unit	
24 hour cumulative dose	
24 hour cumulative dose unit	
◆ Line labelled and checked	Yes
Patent IV access	
◆ Clinician bolus dose given	
Nurse/Midwife Witness	Yes

Note: Refer to the quick reference guide [PCA/NCA – Documenting](#) for instructions on how to document volume infused, changing the rate and changing the bag.

