



ieMR Advanced

PACU Pain Protocol

Quick reference guide

Administration

1. Navigate to the **MAR**. The order will look like this:

	PRN 0.2 mg morphine (Morphine 2.5 mg in sodium chloride 0.9% 5 mL PACU USE ONLY) 0.2 mg, Solution, injectable, IV, every 10 minutes, start: 03/09/18 09:58:00 AEST, PRN for pain, Dose 0.02 mg/kg or 0.04 mg/kg, Indication: - ***PACU USE ONLY*** Dilute 2.5 mg morphine to 5 mL with sodium chlorid... morphine	Not given within 5 days.
--	---	--------------------------

OR this:

	PRN 10 mcg fentanyl (Fentanyl 25 mcg in sodium chloride 0.9% 2.5 mL PACU USE ONLY) 10 mcg, Solution, injectable, IV, every 5 minutes, start: 19-Jul-2018 15:41:00, PRN for pain, Dose 0.2 mcg/kg or 0.5 mcg/kg, Indication: - ***PACU USE ONLY*** Dilute 25 mcg fentanyl to 2.5 mL with sodium chlori... fentanyl	Not given within 5 days.
--	---	--------------------------

2. Perform allergy checks
3. Perform the usual PPID
4. Administer the medication through the MAW
5. Navigate to the **MAR**.
6. Click **Refresh** - the administered dose will now appear in the action cell or on the tile as "Last given:"

	PRN 10 mcg fentanyl (Fentanyl 25 mcg in sodium chloride 0.9% 2.5 mL PACU USE ONLY) 10 mcg, Solution, injectable, IV, every 5 minutes, start: 19-Jul-2018 15:41:00, PRN for pain, Dose 0.2 mcg/kg or 0.5 mcg/kg, Indication: - ***PACU USE ONLY*** Dilute 25 mcg fentanyl to 2.5 mL with sodium chlori... fentanyl	Last given: 10 mcg @ 01-Apr-2019 11:52 AEST	10 mcg Auth
--	---	--	-------------

Documenting Discarded/Unused Opioid in PACU

1. Right click on the green task and select **Chart Details**

	PRN 10 mcg fentanyl (Fentanyl 25 mcg in sodium chloride 0.9% 2.5 mL PACU USE ONLY) 10 mcg, Solution, injectable, IV, every 5 minutes, start: 19-Jul-2018 15:41:00, PRN for pain, Dose 0.2 mcg/kg or 0.5 mcg/kg, Indication: - ***PACU USE ONLY*** Dilute 25 mcg fentanyl to 2.5 mL with sodium chlori... fentanyl	Last given: 10 mcg @ 01-Apr-2019 11:52 AEST	10 mcg Auth	<ul style="list-style-type: none"> Order Info... Chart Details... Quick Chart... Chart Done... Chart Not Done... Unchart... Reschedule This Dose...
--	---	--	-------------	---

In administration window:

1. Enter witness in **Witness by**
2. Tick the box **Not Given** at the bottom of the screen
3. In ***Reason** select **Other-enter comment**
4. Open **Comment** box. Enter amount of opioid discarded. Click **OK**.
5. Click green tick
6. Witness will be prompted for their password




Discontinuing PACU Pain Protocol for Discharging Patients


PACU Pain PowerPlan must be discontinued once the patient is discharged from PACU so it is not available for use outside PACU

To do this:

1. Navigate to the orders screen
2. Rick click on the PowerPlan name **PACU Acute Pain Paediatric** and select **discontinue**





3. In the **Ordering Physician** pop up window:
 - a. Select **Protocol (no co-sign)**
 - b. Enter the prescribing anaesthetists name (found at the top of the order) and click **OK**



4. Click **Orders For Signature** in the bottom right corner

5. Click on the blue circle with the white cross next to the drug order to
6. Enter **Discontinue Reason - No longer required**



7. Click **Sign** in bottom right corner to discontinue the PowerPlan
8. Navigate to the **MAR**, click **Refresh**. Order will now appear in the Discontinued PRN section

